



Agromet Advisory Service Bulletin for Thane District

(Issued jointly by IAAS, Dr. B.S. KonkanKrishiVidyapeeth, & Regional India Meteorological Department, Mumbai)

Dr. Subhash Chavan,
Head,
Department of Agronomy
9422431067

Dr. Vijay More,
Nodal Officer,
Department of Agronomy
9422374001

Prof. Viresh Chavan,
Technical Officer,
Department of Agronomy
9422065344

**Significant past weather for the preceding week
(Period –28/05/2018 to 01/06/2018)**

Rainfall (mm):	0.0
Total Rainfall (mm):	0.0
Total Rainfall (mm) (last year)	2834.2
Maximum temperature (°C)	32.6-37.0
Minimum temperature (°C)	22.6-27.8
Morning RH (%)	76-82
Afternoon RH (%)	57-68
Wind Speed (km/hr):	3.3-5.0
Total cloud cover (Octa)	0

Weather forecast until 08.30 hrs of 06/06/2018

PARAMETERS	Day-1	Day-2	Day-3	Day - 4	Day - 5
	02/06	03/06	04/06	05/06	06/06
Rainfall (mm)	4	5	12	11	17
Maximum temperature (°C)	36	38	35	36	35
Minimum temperature (°C)	29	28	29	29	29
Total cloud cover (Octa)	5	5	6	6	7
Relative Humidity Max. (%)	88	89	87	88	87
Relative Humidity Min. (%)	40	37	44	44	45
Wind Speed (Km/hr)	003	003	002	003	003
Wind Direction	184	248	207	210	201

: Agro-met Advisory:

It is possibility rainfall of 2nd to 6th June, 2018. Sky will cloudy from 2nd to 6th June, 2018. Wind velocity range from 2 to 3 kmph from 2nd to 6th June, 2018.

Crop	Crop Stage	Agromet Advisory
Rice	Field preparation	<ul style="list-style-type: none"> For preparation of rice nursery plough the area & bring the soil to fine tilth and mixed 1 kg. FYM per sq.mt. area. Prepare raised bed of 120 cm. breadth at bottom and 90 cm. on top along the slope of plant. Convenient length of raised bed should be kept according to the slope of land. Apply 1 kg. urea and 3 kg. SSP per guntha at the time of sowing. Store sufficient quantity of improved variety seed and required fertilizers during kharif season.
Sapota	Fruiting Stage	<ul style="list-style-type: none"> If unmatre fruit drops of sapota are observed spray combined fungicides of (Metalxyl-M 8% + Mancozeb 64%) @ 0.2% i.e. 20 g per 10 lit. of water on whole plant canopy. Some parts in the district the sapota fruits are mature size stage, harvest the mature fruits before 10 hours in the morning and after 16 hours in the evening with the help on Atul Sapota harvester. Due to increasing temperature and humidity variation, irrigate the sapota orchard as per the water availability is suggested.
Coconut, Arecanut	--	<ul style="list-style-type: none"> For control of adults and grubs of rhinoceros beetle, dust 2 % Methyl parathion powder after every 2 month interval in FYM pits. To control red palm weevil affected coconut, prepare a slanting hole with the help of screw drill about 1 m height from ground level on tree trunk and pour about 20 ml of 36 % Monocrotophos or 20% Chlorphyriphos with the help of plastic funnel in the hole and close the hole with the help of cement. Clean the arecanut plant by removing dry leaves and other material and spray 1 % bordomixture to control 'koleroga' disease. Select feeding root of arecanut plant then take slant cut to the tip of the root. Prepare 0.3 % solution of Foseryl – Al by mixing 3 gm. per liter. Fill the two plastics bags with 100 ml. above prepare solution and dip the above selected roots in to the plastics bag ensuring the cut portion will always remain in the solution. Tie the bag to the roots. For new plantations of Coconut and Arecanut, clean the area. For coconut dig the pits of 1 x 1 x 1 m. size at the distance of 7.5 x 7.5 m for cashew and arecanut pits of 0.6 x 0.6 x 0.6 m. size at the distance of 2.7 x 2.7 m re-filled the pits with mixture of soil, 20 to 30 kg of FYM or Compost and 2 kg single super phosphate.
Vegetable and Fruit crop nursery	--	<ul style="list-style-type: none"> Start sowing of cucurbitaceous crops with spacing of 1.5 m x 90 cm, at the time of sowing applies 2 kg FYM or compost alongwith 50 to 60 g mix fertilizer (Suphala). Start sowing of okra in ridges and furrow having spacing of 60 x 45 cm. Store sufficient quantity of F.Y.M. and loam soil required for polythene bag filling during rainy season for raising the seedling or grafts.
Goats/ Milch animals/ Poultry	--	<ul style="list-style-type: none"> Provide clean and hygienic drinking water and nutritious fodder and concentrates to farm animal. Provide 1 to 1.5 kg concentrate + 15 to 20 kg green fodder + 4 to 5 kg roughages per day for milch animals. Young goats provide 3 to 4 kg green fodder + 2 to 2.5 kg dry fodder + 300 g concentrate,. Protect the animals from afternoon heat. Keep good aeration in animal shed. For boiler poultry bird 1st three week provide boiler starter and 4 to 6 week old bird provide boiler finisher as per their daily requirement.
Suggestion	--	<ul style="list-style-type: none"> Contact nearby SAU Scientists or State Agril. Dept. for detail control measures against incidence of pest and diseases under adverse weather conditions.