



Agromet Advisory Service Bulletin for Ratnagiri District
(Issued jointly by IAAS, Dr. B.S. Konkan Krishi Vidyapeeth,
& Regional India Meteorological Department, Mumbai)
(02358) 282387



No. 05/2019

Date: 15/01/2019

Duration – 5 days

Dr. Subhash Chavan,
Head,
Department of Agronomy
9422431067

Dr. Vijay More,
Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav,
Technical Officer,
Department of Agronomy
8379901160

Significant past weather for the preceding week (Period –09/01/2019 to 15/01/2019)							Weather Parameters	Weather forecast until 08.30 hrs of 20/01/2019				
09/01	10/01	11/01	12/01	13/01	14/01	15/01		16/01	17/01	18/01	19/01	20/01
0.0	0.0	0.0	0.0	0.0	0.0	0.0	Rainfall (mm)	0	0	0	0	0
32.6	32.2	33.0	31.0	30.2	32.0	33.0	Maximum temperature (°C)	32	32	33	33	33
10.5	11.2	12.0	11.0	10.0	11.5	14.0	Minimum temperature (°C)	17	17	17	17	17
0	0	0	0	0	0	0	Cloud cover (Octa)	0	0	0	0	1
88	85	88	85	88	85	91	Relative Humidity Max. (%)	60	60	58	58	58
62	60	57	59	58	60	55	Relative Humidity Min. (%)	35	35	35	35	35
-	-	-	-	-	-	-	Wind speed (Km/hr)	9	9	9	9	7
calm	calm	calm	calm	calm	calm	calm	Wind direction	64	46	49	63	60
Rainfall (mm) in last week						Rainfall (mm) from 01/01/2019 to till dated		Total Rainfall (mm) in last year				
0.0						0.0		3071.8				

Agro-met Advisory

There will be normal weather condition from 16th to 20th January, 2019.

Crop	Stage	Agro Advise
Lablab bean	Flowering	<ul style="list-style-type: none"> Provide irrigation to lablab bean crop when crop is in flowering and pod filling stage.
Mango	Flowering	<ul style="list-style-type: none"> As per blossom protection schedule for mango crop, spray Imidacloprid 17.8% SL @ 6 ml per 10 liter of water to control the incidence of mango hoppers and midge fly before the flower opening to avoid the adverse effect on pollinators. Also add Hexaconazole 5% @ 5 ml or wettable Sulphur 80% @ 20 gm per 10 liter in water for control of powdery mildew disease.
Cashewnut	Fruiting	<ul style="list-style-type: none"> There is possibility of incidence of tea mosquito bug and thrips during fruit bearing stage of cashewnut, if incidence is noticed spray Lambda cyhalothrin 5%EC @6 ml per 10 liter of water. (insecticide is not under label claim) If the incidence of Cashew Stem and Root Borer (CSRB) is noticed in Cashew, remove the grubs from the holes with the help of 15 mm chisel and apply Chloropyriphos 20 EC @ 5 ml per liter of water on the stem (swabbing) and pour remaining solution around the stem near to the roots.
Vegetables/ Fruit crop nursery	Fruiting	<ul style="list-style-type: none"> Install cue lure 'Rakshak' trap @ 4nos. per hectare at the time of initiation of in watermelon crop for effective control of fruit fly. Provide irrigation to fruit crop nursery regularly.
<p>This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.</p>		