



**Agromet Advisory Service Bulletin for Raigad District**  
(Issued jointly by IAAS, Dr. B.S. Konkan Krishi Vidyapeeth,  
& Regional India Meteorological Department, Mumbai)  
(02358) 282387



No. 08/2019

Date: 25/01/2019

Duration – 5 days

**Dr. Subhash Chavan,**  
Head,  
Department of Agronomy  
9422431067

**Dr. Vijay More,**  
Nodal Officer,  
Department of Agronomy  
9422374001

**Dr. Shital Yadav,**  
Technical Officer,  
Department of Agronomy  
8379901160

Significant past weather for the preceding week (Period –19/01/2019 to 25/01/2019)							Weather Parameters	Weather forecast until 08.30 hrs of 30/01/2019				
19/01	20/01	21/01	22/01	23/01	24/01	25/01		26/01	27/01	28/01	29/01	30/01
0.0	0.0	0.0	0.0	-	-	-	Rainfall (mm)	0	0	0	0	0
37.0	36.4	36.0	33.2	-	-	-	Maximum temperature (°C)	30	31	31	32	32
13.8	14.7	14.8	12.6	-	-	-	Minimum temperature (°C)	17	17	17	18	18
0	0	0	0	-	-	-	Cloud cover (Octa)	0	0	1	1	0
89.0	89.0	87.0	86.0	-	-	-	Relative Humidity Max.(%)	74	72	72	70	70
25.0	25.0	29.0	-	-	-	-	Relative Humidity Min. (%)	71	71	71	69	69
2.2	1.9	2.8	-	-	-	-	Wind speed (Km/hr)	13	14	16	14	10
Calm	E	Calm	-	-	-	-	Wind direction	22	44	59	63	70
Rainfall (mm) in last week						Rainfall (mm) from 01/01/2019 to till dated			Total Rainfall (mm) in last year			
0.0						0.0			3558.3			

**Agro-met Advisory**  
There will be normal weather condition from 26<sup>th</sup> to 30<sup>th</sup> January, 2019. Wind velocity range from 10 to 16 Kmph.

Crop	Stage	Agro Advise
Rice	Seedling	<ul style="list-style-type: none"> <li>In rice field, maintain optimum water level of 2.5- 5 cm upto 30 days after transplanting.</li> </ul>
Lablab bean	Flowering and pod filling stage	<ul style="list-style-type: none"> <li>There is possibility for incidence of pod borer on lablab bean, if incidence is noticed, collect and destroy all infected pods and spray Dimethoate 30% EC @ 20 ml per 10 liter of water.</li> </ul>
Mango	Flowering	<ul style="list-style-type: none"> <li>Provide support to newly planted mango graft, as there is forecast of high wind speed.</li> <li>Low temperature during night affects the flower initiation in mango. For inducing flowering in mango, burned the dried leaves and create smoke under the canopy of the mango tree or spray mango tree with 1% potassium nitrate solution (1 kg KNO<sub>3</sub> per 100 liter of water)</li> </ul>
Cashewnut	Fruiting	<ul style="list-style-type: none"> <li>Provide support to newly planted cashew graft, as there is forecast of high wind speed.</li> <li>There is possibility of incidence of tea mosquito bug and thrips during fruit bearing stage of cashewnut, if incidence is noticed spray Lambda cyhalothrin 5%EC @6 ml per 10 liter of water. ( insecticide is not under label claim)</li> </ul>
Vegetables/ Fruit crop nursery	Fruiting	<ul style="list-style-type: none"> <li>Due to forecast of high windspeed follow earthing up operation in vegetable crops.</li> <li>There is possibility for incidence of aphids on watermelon, if incidence is noticed spray Dimethoate @10 ml or Malathion @20 ml per 10 liter of water. Also use yellow stick cards in field.</li> <li>Provide irrigation to fruit crop nursery, vegetable crops and newly planted plantation crops at evening hours.</li> </ul>
Milch animal	-	<ul style="list-style-type: none"> <li>Protect the animals /poultry birds from low temperature by providing curtains and electric bulbs as per the need.</li> <li>Provide clean and hygienic drinking water to farm animals.</li> <li>For control of worms in goat, give Albendazol 7.5 ml per kg weight of goat.</li> </ul>

**This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of “Gramin Krishi Mausam Sewa (GKMS)” Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli.**  
For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.