



Agromet Advisory Service Bulletin for Raigad District
(Issued jointly by IAAS, Dr. B.S. Konkarn Krishi Vidyapeeth,
& Regional India Meteorological Department, Mumbai)
(02358) 282387



No. 54/2019

Date: 05/07/2019

Duration – 5 days

Dr. Ashokkumar Chavan,
Head,
Department of Agronomy
9422373396

Dr. Vijay More,
Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav,
Technical Officer,
Department of Agronomy
8379901160

Significant past weather for the preceding week (Period –29/06/2019 to 05/07/2019)							Weather Parameters	Weather forecast until 08.30 hrs of 10/07/2019				
29/06	30/06	01/07	02/07	03/07	04/07	05/07		06/07	07/07	08/07	09/07	10/07
287.2	106.8	83.4	136.6	4.8	13.2	35.4	Rainfall (mm)	50	70	65	59	75
-	-	-	-	-	-	-	Maximum temperature (°C)	30	29	30	30	29
-	-	-	-	-	-	-	Minimum temperature (°C)	27	27	25	25	25
-	-	-	-	-	-	-	Cloud cover (Octa)	8	8	7	8	8
-	-	-	-	-	-	-	Relative Humidity Max.(%)	89	91	91	90	92
-	-	-	-	-	-	-	Relative Humidity Min. (%)	81	80	81	81	81
-	-	-	-	-	-	-	Wind speed (Km/hr)	18	15	16	21	18
-	-	-	-	-	-	-	Wind direction	227	225	228	236	246
Rainfall (mm) in last week						Rainfall (mm) from 01/01/2019 to till dated		Total Rainfall (mm) in last year				
667.4						807.4		3558.3				

Agro-met Advisory

There is possibility of heavy rainfall from 6th to 10th July, 2019 and wind velocity ranges from 16 to 21 Kmph.
Extended range rainfall forecast for Konkarn division for the period of 5th to 11th July 2019 is above normal.

Crop	Stage	Agro Advise
Kharif rice	Seedling	<ul style="list-style-type: none"> Repair the bunds of rice field and keep bunds weed free. Due to forecast of heavy rain, make arrangement for drain out excess water from rice nursery and maintain optimum water level of 2.5 to 5cm upto 30 days after transplanting. Prepare field for transplanting by puddling. At the time of puddling apply 87kg urea, 313 kg single super phosphate and 84 kg muriate of potash per hectare. If incorporation of 5 tonnes of glyricidia leaves per hectare is done then reduce the dose of nitrogenous fertilizers to 50 percent. <p>Note: Apply fertilizer during puddling when intensity of rainfall is low</p> <ul style="list-style-type: none"> Rice seedling with 5 to 6 leaves with height of 12 to 15 cm and 20 to 27 days old are consider ideal for transplanting. It is recommended to transplant 20 days old seedling in case of early rice varieties and 27 to 30 days old seedling in case of midlate and late rice varieties. Transplant 2-3 rice seedling per hill in case of straight varieties and one seedling per hill in case of hybrid at a spacing of 15 x 15 cm (early) or 20 x 15 cm (midlate and late varieties). follow upright and shallow (2.5 to 3.5 cm) transplanting of seedling. For control of crabs from rice field, use bait made up of 1 kg of cook rice and 75 gm of insecticide Acephate 75WP. Prepare 100 arecanut size balls and keep them in each crab burrow and close it with wet mud. On the next day if some of the burrows are found open, again use the same bait.
Finger millet	Seedling	<ul style="list-style-type: none"> Due to forecast of heavy rain, make arrangement for drain out excess water from nursery. Carryout weeding in finger millet nursery. For preparation of field finger millet, carryout ploughing and incorporate 5 tonnes of FYM or compost per hectare at the time of ploughing.
New plantation	Vegetative	<ul style="list-style-type: none"> Due to forecast for increase in wind speed, provide support to newly planted fruit crops.
Coconut/ Arecanut	-	<ul style="list-style-type: none"> Due to forecast of heavy rain, make arrangement for drain out excess water from coconut and arecanut orchard
Vegetable crops	Vegetative	<ul style="list-style-type: none"> Due to forecast of heavy rain, make arrangement for drain out excess water from vegetable crop field. Proper support should be given to cucurbitaceous crops.
Milch animals/Goat /Poultry	-	<ul style="list-style-type: none"> Provide clean and hygienic drinking water to farm animals also maintain good aeration in farm shed. Protect the farm animals from rain water. Carryout planting of biennial and perennial forage crops for animal. The varieties of perennial crops (stumps) like paragrass, Hybrid napier (Jayawant CO- 3, Yashwant) etc.

This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkarn Krishi Vidyapeeth, Dapoli.
For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.

