

**Agromet Advisory Service Bulletin for Raigad District**
(Issued jointly by IAAS, Dr. B.S. KonkanKrishiVidyapeeth, & Regional India Meteorological Department, Mumbai)**Dr. Subhash Chavan,**
Head,
Department of Agronomy
9422431067**Dr. D.N. Jagtap,**
Nodal Officer,
Department of Agronomy
9403988143**Prof. Viresh Chavan,**
Technical Officer,
Department of Agronomy
9422065344**Significant past weather for the preceding week**
(Period –02/02/2018 to 06/02/2018)

Rainfall (mm):	
Total Rainfall (mm):	
Total Rainfall (mm) (last year)	
Maximum temperature (°C)	
Minimum temperature (°C)	
Morning RH (%)	
Afternoon RH (%)	
Wind Speed (km/hr):	
Total cloud cover (Octa)	

Weather forecast until 08.30 hrs of 11/02/2018

PARAMETERS	Day-1	Day-2	Day-3	Day - 4	Day - 5
	07/02	08/02	09/02	10/02	11/02
Rainfall (mm)	0	0	0	0	0
Maximum temperature (°C)	29	29	29	28	28
Minimum temperature (°C)	21	20	19	19	20
Total cloud cover (Octa)	8	2	4	7	7
Relative Humidity Max. (%)	60	60	71	72	71
Relative Humidity Min. (%)	40	41	43	52	46
Wind Speed (Km/hr)	005	004	004	004	002
Wind Direction	126	70	118	22	116

: Agro-met Advisory:**Sky will be cloudy from 7th to 11th February, 2018. Wind velocity range from 2 to 5 kmph from 7th to 11th February, 2018.**

Crop	Crop Stage	Agromet Advisory
Summer Rice	Seedling	<ul style="list-style-type: none"> Transplant paddy seedling after 30 to 35 days sowing with spacing 20 x 15 cm. Before transplanting apply 130 kg urea + 312.5 kg single super phosphate + 83 kg murate of potash per ha.
Pulses crops	Vegetative stage	<ul style="list-style-type: none"> There are forecasted cloudy weather on 7th to 11th February, 2018 also variation between minimum temperature hence incidence of aphids and leaf minor is increased if it is observed on pulse crops, spray 10 ml Dimethoate or Lambda cyhalothrin 5% EC @ 6ml. per 10 lit. of water at an interval of 15-20 days.
Mango and Cashew	Plantation	<ul style="list-style-type: none"> There are forecasted cloudy weather for 7th to 11th February, hence possibility of incidence of mango hopper, if it is noticed spray Imidachloprid 17.8% SL. @ of 3 ml per 10 liter of water for management of powdery mildew mix Hexaconazole 5% EC @ 5 ml or Sulphur 80% WP @ 20 gm per 10 lit of water. There are forecasted cloudy weather for 7th to 11th February, 2018 hence possibility of incidence of Tea mosquito on new emerging flush of cashew is noticed, spray Lambda Cyhalothrin 5% EC (6 ml. in 10 lit. of water). (These insecticide are not under label claim)
Coconut, Arecanut	--	<ul style="list-style-type: none"> For control of adults and grubs of rhinoceros beetle, dust 2 % Methyl parathion powder after every 2 month interval in FYM pits. To control red palm weevil affected coconut, prepare a slanting hole with the help of screw drill about 1 m height from ground level on tree trunk and pour about 20 ml of 36 % Monocrotophos or 20% Chlorphyriphos with the help of plastic funnel in the hole and close the hole with the help of cement.
Vegetable crop and watermelon crop	--	<ul style="list-style-type: none"> For control of bacterial wilt of chilli, brinjal and tomato seedling, spray Copper Oxychloride @ 20 to 25 g/10 lit of water on seedbed after sowing of seeds. Also apply Carbarly or Lindane powder on side of seedbed to control against ant. There is possibility for incidence of powdery mildew, spray Hexaconazole 5% EC 5 ml or Sulphur 80% WP @ 20 g per 10 lit. of waters. Due to cloudy weather on 7th to 11th February, 2018 there is possibility for increased incidence of hopper, aphids and thrips on vegetable crops viz., brinjal, tomato, cabbage, chilli, knol knol etc., if it is noticed spray Malathion @ 20 ml or Dimethoate @ 12 ml per 10 liter of water. Cucurbitaceous vegetables are in fruit bearing stage, install of Rakshak fruit fly traps developed by University @ 4 traps per ha area is advocated to control fruit fly in vegetables garden. In watermelon crop if symptoms of wilting are observed then drenching of Carbandazim @ 1 gm. per lit. of water per vine. Apply straw mulch or polythene mulch for water melon crops.
Goats/ Milch animals	--	<ul style="list-style-type: none"> Provide clean and hygienic drinking water and nutritious fodder and concentrates to farm animal. Protect the animals/poultry birds from low temperature by providing curtains and electric bulbs as per the need.
Suggestion	--	<ul style="list-style-type: none"> Contact nearby SAU Scientists or State Agril. Dept. for detail control measures against incidence of pest and diseases under adverse weather conditions.