



**Agromet Advisory Service Bulletin for Raigad District**  
(Issued jointly by IAAS, Dr. B.S. KonkanKrishiVidyapeeth, & Regional India Meteorological Department, Mumbai)

**Dr. Subhash Chavan,**  
Head,  
Department of Agronomy  
9422431067

**Dr. D.N. Jagtap,**  
Nodal Officer,  
Department of Agronomy  
9403988143

**Prof. Viresh Chavan,**  
Technical Officer,  
Department of Agronomy  
9422065344

**Significant past weather for the preceding week**  
(Period –04/12/2016 to 08/12/2017)

Rainfall (mm):	
Total Rainfall (mm):	
Total Rainfall (mm) (last year)	
Maximum temperature (°C)	
Minimum temperature (°C)	
0Morning RH (%)	
53Afternoon RH (%)	
Wind Speed (km/hr):	
Total cloud cover (Octa)	

**Weather forecast until 08.30 hrs of 13/12/2017**

PARAMETERS	Day-1	Day-2	Day-3	Day - 4	Day - 5
	09/12	10/12	11/12	12/12	13/12
Rainfall (mm)	0	0	0	0	0
Maximum temperature (°C)	30	30	31	30	29
Minimum temperature (°C)	21	21	19	19	18
Total cloud cover (Octa)	1	0	0	0	0
Relative Humidity Max. (%)	90	73	77	88	90
Relative Humidity Min. (%)	48	33	34	40	48
Wind Speed (Km/hr)	005	005	003	003	003
Wind Direction	78	82	57	46	57

**: Agro-met Advisory:**

Sky will be slightly cloudy on 9<sup>th</sup> December, 2017. Wind velocity range from 3 to 5 kmph from 9<sup>th</sup> to 13<sup>th</sup> December, 2017.

Crop	Crop Stage	Agromet Advisory
<b>Mango and Cashew</b>	<b>Plantation</b>	<ul style="list-style-type: none"> <li>If incidence of mango hopper is noticed spray Lambda Cyhalothrin @5% E.C. @ of 6 ml per 10 liter of water for management of Anthracnose mix Carbendazim 12% + Mancozeb 63% @ 10 g per 10 lit of water.</li> <li>If incidence of Tea mosquito on new emerging flush of cashew is noticed, spray profenophos 15% (10 ml. in 10 lit. of water). (These insecticide are not under label claim)</li> </ul>
<b>Coconut, Arecanut</b>	--	<ul style="list-style-type: none"> <li>For controlling of bud rot of coconut palm, clean infected bud and drenching with 1% Bordo mixture or copper-oxy-chloride @ 25 g per 10 lit of water.</li> <li>Spraying the FYM pits with 0.2 % Carbaryl for control of adults and grubs of rhinoceros beetle.</li> <li>To control red palm weevil affected coconut, fill the holes made by RPW with 10 per cent Carbaryl dust and sand mixture. Prepare a slanting hole with the help of screw drill about 1 m height from ground level on tree trunk and pour about 20 ml of 36 % Monocrotophos or 20% Chlorphyriphos with the help of plastic funnel in the hole and close the hole with the help of cement.</li> </ul>
<b>Vegetable crop and watermelon crop</b>	--	<ul style="list-style-type: none"> <li>For control of bacterial wilt of chilli, brinjal and tomato seedling, spray Copper Oxychloride @ 20 to 25 g/10 lit of water on seedbed after sowing of seeds. Also apply Carbarly or Lindane powder on side of seedbed to control against ant.</li> <li>There is possibility for incidence of powdery mildew, spray Hexaconazole 5% EC 5 ml or Sulphur 80% WP @ 20 g per 10 lit. of waters.</li> <li>For cultivation of watermelon ploughing and cloud crushing of land and furrow be prepared at 4 m length 90 cm apart the both side of furrow prepared 30 x 30 x30 cm size pit, apply 1 to 1.5 kg FYM + 10 gm Carbaryl (10%) Powder, mix thoroughly with soil, sowing of 3 to 4 seeds at the depth of 2 to 2.5 cm, thinning after 15 days emergence of seedling keep only 2 seedlings per pit. The 50 kg N + 50 kg P<sub>2</sub>O<sub>5</sub> and 50 kg K<sub>2</sub>O per ha was required. The 1/3<sup>rd</sup> dose of N and full dose of P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O be applied at the time of sowing and remaining N be apply 1 and 2 months after sowing at equal quantity.</li> </ul>
<b>Goats/ Milch animals</b>	--	<ul style="list-style-type: none"> <li>Provide clean and hygienic drinking water and nutritious fodder and concentrates to farm animal.</li> <li>Protect the animals/poultry birds from low temperature by providing curtains and electric bulbs as per the need.</li> </ul>
<b>Suggestion</b>	--	<ul style="list-style-type: none"> <li>Contact nearby SAU Scientists or State Agril. Dept. for detail control measures against incidence of pest and diseases under adverse weather conditions.</li> </ul>