



Agromet Advisory Service Bulletin for Thane District
(Issued jointly by IAAS, Dr. B.S. KonkanKrishiVidyapeeth, & Regional India Meteorological Department, Mumbai)

Dr. Uttam Mahadkar,
Head,
Department of Agronomy
9422791998

Dr. D.N. Jagtap,
Nodal Officer,
Department of Agronomy
9403988143

Prof. Viresh Chavan,
Technical Officer,
Department of Agronomy
9422065344

Significant past weather for the preceding week
(Period -07/04/2017 to 11/04/2017)

Rainfall (mm):	
Total Rainfall (mm):	
Total Rainfall (mm) (last year)	
Maximum temperature (°C)	
Minimum temperature (°C)	
Morning RH (%)	
Afternoon RH (%)	
Wind Speed (km/hr):	
Total cloud cover (Octa)	

Weather forecast until 08.30 hrs of 16/04/2017

PARAMETERS	Day-1	Day-2	Day-3	Day - 4	Day - 5
	12/04	13/04	14/04	15/04	16/04
Rainfall (mm)	0	0	0	0	0
Maximum temperature (°C)	36	36	35	35	34
Minimum temperature (°C)	25	24	23	22	22
Total cloud cover (Octa)	0	0	0	0	0
Relative Humidity Max. (%)	33	34	61	67	67
Relative Humidity Min. (%)	16	14	15	23	24
Wind Speed (Km/hr)	003	005	006	006	005
Wind Direction	83	187	180	184	177

: Agro-met Advisory:

Sky will be clear from 12th to 16th April, 2017. Wind velocity range from 3 to 6 kmph on 12th to 16th April, 2017.

Crop	Crop Stage	Agromet Advisory
Sapota	Vegetative Stage	<ul style="list-style-type: none"> If unripe fruit drops of sapota are observed spray combined fungicides of (Metalxyl-M 8% + Mancozeb 64%) @ 0.2% i.e. 20 g per 10 lit. of water on whole plant canopy. Some parts in the district the sapota fruits are mature size stage, harvest the mature fruits before 10 hours in the morning and after 16 hours in the evening with the help on Atul Sapota harvester.
Coconut, Arecanut	--	<ul style="list-style-type: none"> Spraying the FYM pits with 0.2 % Carbaryl for control of adults and grubs of rhinoceros beetle. To control red palm weevil affected coconut, fill the holes made by RPW with 10 per cent Carbaryl dust and sand mixture. Prepare a slanting hole with the help of screw drill about 1 m height from ground level on tree trunk and pour about 20 ml of 36 % Monocrotophos or 20% Chlorpyrifos + 20 ml water with the help of plastic funnel in the hole and close the hole with the help of cement.
Vegetable	Vegetative Stage	<ul style="list-style-type: none"> If incidence of hopper, aphids and thrips is observed in <i>rabi</i> vegetable crops viz., brinjal, tomato, cabbage, chilli, knol knol etc., spray Malathion @ 20 ml or Dimethoate @ 12 ml per 10 liter of water. There is possibility of incidence of fruit fly on cucurbitaceous crops, install 'Rakshak' fruit fly traps developed by University @ 4 traps per ha. area.
Watermelon Cultivation	--	<ul style="list-style-type: none"> Harvest the mature fruits of watermelon by observing weathered tendril near peduncle of fruits and after completion of duration of variety. Installation of Rakshak fruit fly traps developed by the University @ 4 traps per ha area is advocated to control fruit fly in cucurbitaceous crops as well as watermelon crop.
Goats/ Milch animals	--	<ul style="list-style-type: none"> Provide clean and hygienic drinking water and nutritious fodder and concentrates to farm animal. Planting of binneal forage crops like Paragrass, Napier, Gajraj etc is advocated.
Suggestion	--	<ul style="list-style-type: none"> Contact nearby SAU Scientists or State Agril. Dept. for detail control measures against incidence of pest and diseases under adverse weather conditions.