



Agromet Advisory Service Bulletin for Raigad District
(Issued jointly by IAAS, Dr. B.S. KonkanKrishiVidyapeeth, & Regional India Meteorological Department, Mumbai)

Dr. Subhash Chavan,
Head,
Department of Agronomy
9422431067

Dr. D.N. Jagtap,
Nodal Officer,
Department of Agronomy
9403988143

Prof. Viresh Chavan,
Technical Officer,
Department of Agronomy
9422065344

Significant past weather for the preceding week
(Period –08/09/2016 to 12/09/2017)

Rainfall (mm):	
Total Rainfall (mm):	
Total Rainfall (mm) (last year)	
Maximum temperature (°C)	
Minimum temperature (°C)	
Morning RH (%)	
Afternoon RH (%)	
Wind Speed (km/hr):	
Total cloud cover (Octa)	

Weather forecast until 08.30 hrs of 17/09/2017

PARAMETERS	Day-1	Day-2	Day-3	Day - 4	Day - 5
	13/09	14/0809	15/09	16/09	17/09
Rainfall (mm)	5	15	20	6	0
Maximum temperature (°C)	33	33	33	32	32
Minimum temperature (°C)	26	26	25	25	25
Total cloud cover (Octa)	8	7	8	8	6
Relative Humidity Max. (%)	87	84	85	86	84
Relative Humidity Min. (%)	77	75	69	73	72
Wind Speed (Km/hr)	002	004	007	007	005
Wind Direction	164	117	158	169	139

: Agro-met Advisory:

It is possibility rainfall of 13th to 16th September, 2017. Sky will cloudy from 13th to 17th September, 2017. Wind velocity range from 2 to 7 kmph from 13th to 17th September, 2017.

Crop	Crop Stage	Agromet Advisory
Rice	Panicle initiation	<ul style="list-style-type: none"> Farmers are requested to supervise daily rice transplanted field to observe pests and diseases. Maintain water level 5 cm by bunding rice field. Apply 3rd split dose of fertilizer to the mid-late and late rice varieties at the time of flowering of rice crop @ 20 Kg N/ha. There is a possibility for incidence of blue beetle, if it is noticed; spray Quinalphos 25% EC @ 2 lit. or or Triazophos 40% EC @ 625 ml or lambdacyhalothrin 5% EC @ 250 ml in 500 lit. of water per hecter. There is a possibility for incidence of fungal disease of blast on rice crop, if incidence is noticed; spray any one fungicides of Tricyclazole (75% WP) @ 10 gm per 10 lit of water or Isoprothiolane 40% @ 10 ml per 10 lit or water.
Mango and Cashew	Plantation	<ul style="list-style-type: none"> Weeding should be followed in Mango and Cashew garden with the help of sickle/grass cutting machine or spray Glyphosate weedicide @ 3-4 ml per lit water for effective control of weeds. <p>Note : To avoid grazing of animal on weedicide spray in mango and cashew garden.</p>
Coconut, Arecanut	--	<ul style="list-style-type: none"> Spraying the FYM pits with 0.2 % Carbaryl for control of adults and grubs of rhinoceros beetle. To control red palm weevil affected coconut, fill the holes made by RPW with 10 per cent Carbaryl dust and sand mixture. Prepare a slanting hole with the help of screw drill about 1 m height from ground level on tree trunk and pour about 20 ml of 36 % Monocrotophos or 20% Chlorpyrifos with the help of plastic funnel in the hole and close the hole with the help of cement. Apply osety AL 0.3% (30 g/10 lit. water) + Urea Suphala 111 briquettes or Fosety AL 0.3% (30 g/10 lit. water) + Annapurana 76 briquettes per arecanut palm to avoid fungal immature fruit drop in coconut and Koleroga disease (immature nut drop) in arecanut.
Vegetable crop	Fruiting stage	<ul style="list-style-type: none"> Install methyl ugenol “Rakshak” tarp @ 4 nos. per ha in cucurbitaceous vegetable crops, for effective control of fruit fly.
Goats/ Milch animals	--	<ul style="list-style-type: none"> Provide clean and hygienic drinking water and nutritious fodder and concentrates to farm animal. Protect the poultry birds from low temperature by providing electric bulbs as per the need.
Suggestion	--	<ul style="list-style-type: none"> Contact nearby SAU Scientists or State Agril. Dept. for detail control measures against incidence of pest and diseases under adverse weather conditions.