



Agromet Advisory Service Bulletin for Raigad District
(Issued jointly by GKMS, Dr. B.S. Konkan Krishi Vidyapeeth,
& Regional India Meteorological Department, Mumbai)
(02358) 282387



No. 38/2020

Date: 12/05/2020

Duration – 5 days

Dr. Prashant Bodake,
Head,
Department of Agronomy
9420413255

Dr. Vijay More,
Nodal Officer,
Department of Agronomy
9422374001

Dr. Shital Yadav,
Technical Officer,
Department of Agronomy
8379901160

Significant past weather for the preceding week (Period –06/05/2020 to 12/05/2020)							Weather Parameters	Weather forecast until 08.30 hrs of 17/05/2020				
06/05	07/05	08/05	09/05	10/05	11/05	12/05		13/05	14/05	15/05	16/05	17/05
-	-	-	-	-	-	-	Rainfall (mm)	0	0	5	5	3
-	-	-	-	-	-	-	Maximum temperature (°C)	33	33	32	33	33
-	-	-	-	-	-	-	Minimum temperature (°C)	26	26	26	27	27
-	-	-	-	-	-	-	Cloud cover (Octa)	1	2	3	1	3
-	-	-	-	-	-	-	Relative Humidity Max. (%)	68	76	76	77	86
-	-	-	-	-	-	-	Relative Humidity Min. (%)	28	33	33	38	47
-	-	-	-	-	-	-	Wind speed (Km/hr)	4	5	5	5	5
-	-	-	-	-	-	-	Wind direction	SSW	WNW	SE	WSW	W
Rainfall (mm) in last week							Rainfall (mm) from 01/01/2020 to till dated	Total Rainfall (mm) in last year				
0.0							0.0	5197.2				

Agro-met Advisory

There is possibility of light rainfall from 15th to 17th May 2020 at isolated places and also possibility of increase in maximum and minimum temperature and sky remain partly cloudy.

Crop	Stage	Agro Advise
Precaution to be taken by farmers due to Covid-19		<ul style="list-style-type: none"> In order to avoid the field level contagious spread of Corona virus, farmers are advised to follow general precautions and safety measures in harvesting, post-harvest operations, storage and marketing of produce
Value addition of Fruits	--	<ul style="list-style-type: none"> Due to the lockdown in the view of outbreak of Covid-19 (Corona), farmers are facing difficulties in marketing of perishable farm produce. To reduce the loss of perishable fruits, it is suggested to go for value addition by processing the fruits. In this connection it is possible to make pickle and chutney from unripe mango fruits, pulp and leather from ripe mango, fried chips and leather from jackfruits, syrup from available cashew apples, pickle, chutney, and syrup from Karonda fruits, amsul, syrup and agal from ripe kokum, syrup from jamun fruits. Detail flow chart for preparation of vale added product is available in farmers corner on university website (www.dbskkv.org).
Summer rice	Maturity	<ul style="list-style-type: none"> Due to forecast for light rain from 15th to 17th May,2020, harvest the matured rice varieties in morning hours near to ground level and follow immediate threshing, threshed grains should be stored in protected shed.
Kharif rice	Land preparation	<ul style="list-style-type: none"> Start the land preparation operation for kharif season and expose soil to sun which will help in controlling weeds effectively. Procure the rice seed as per requirement for kharif season. Ratnagiri-1, Ratnagiri-2, Ratnagiri-3, Ratnagiri-7, Karjat-3 and Karjat -5 for bold seeded, Ratnagiri-4, Ratnagiri-5, Ratnagiri-6, Ratnagiri-24, Phondaghat-1 also Sahyadri, Sahyadri-2. Sahyadri-3, Sahyadri-4 and Sahyadri-5 hybrid varieties, for fine seeded, Karjat-4 is for very fine seeded and Panvel-1, Panvel-2 and Panvel-3 for Kharlands categories are recommended by Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli for cultivation in Konkan region during Kharif. Recommended seed rate for paddy is 20 to 24 kg acre⁻¹ for bold varieties, 14 to 16 kg acre⁻¹ for fine varieties and 8 kg acre⁻¹ for hybrid varieties. For availability of seeds and planting materials, please contact to University seed department tel.no. 8275013396 during office hours.
Mango	Fruiting	<ul style="list-style-type: none"> Due to forecast for light rain from 15th to 17th May,2020, harvest the mature mango fruits before 10 hours in the morning with the help on Nutan mango harvester at 80 to 85% maturity. Keep the harvested fruits in shade to prevent development of spongy tissue disease and protect from heat. Do not spray any insecticides/fungicides 15 days before harvesting of mango fruits. To prevent incidence of post-harvest diseases such as fruit rot on mango fruits, place the fruits in hot water of 52°C for 10 minutes and then keep for ripening. Use C.F.B. (corrugated fiber box) for packing developed by B. S. Konkan Krishi Vidyapeeth, Dapoli. Transport of harvested fruits should be done preferably during night hours. Collect and destroy all fruit fly and fruit borer infected fallen fruits regularly and keep mango orchard clean.

		<ul style="list-style-type: none"> • Due to increase in rate of evaporation, provide irrigation to newly planted mango orchard. • There is possibility of incidence of Stem borer in mango tree. Grubs tunnels in the sapwood on the trunk and branches and results into wilting of branches or entire tree. Observe the mango orchard regularly for pest infestation. If the incidence is noticed, remove the grubs from the holes and pour solution of Chloropyriphos 20 EC @ 10 ml + 50 ml Kerosene into hole and seal the hole.
Mango High density planting	--	<ul style="list-style-type: none"> • After complete harvest of mango fruits from high density (5 X 5 m or 6 X 4 m) orchard, go for light pruning. It includes detopping, pruning of cross branches and removing of dead wood. The height of high-density orchard should be maintained at 80% of row distance.
Cashewnut	Fruiting	<ul style="list-style-type: none"> • Due to increase in rate of evaporation, provide irrigation to newly planted cashewnut orchard. • Clean cashewnut orchard by removing all diseased and dried branches to maintain good sanitation in orchard. Apply bordopaste on cut portion of tree.
Coconut	--	<ul style="list-style-type: none"> • Due to forecast for increase in temperature and decrease in humidity, provide irrigation to coconut orchard at 5-6 days interval also use straw mulch to reduce evaporation losses.
Arecanut	--	<ul style="list-style-type: none"> • Due to forecast for increase in temperature and decrease in humidity, provide irrigation to arecanut orchard at 4-5 days interval also use straw mulch to reduce evaporation losses. • Clean the arecanut orchard by removing dry leaves and infected inflorescence to maintain good sanitation in orchard.
Vegetable crops	Sowing	<ul style="list-style-type: none"> • If irrigation facility is available then go for sowing nursery of brinjal, chilli and tomato vegetables crops for kharif season. Prepare raised bed of 3 m length x 1 m breadth x 15 cm height. Apply 5 kg FYM, 35 gm urea, 100 gm single super phosphate and 25 gm muriate of potash per sq. m. Treat the seed with thiram fungicide @ 3 gm/kg of seed before sowing. To protect the seedling from wilt disease, drenching of 1% bordomixture 3 to 4 days before sowing on nursery bed is suggested.
Fruit crop nursery		<ul style="list-style-type: none"> • There is forecast for increase in temperature and decrease in afternoon relative humidity, provide sufficient irrigation to fruit crop nurseries also provide shed.
Milch animals	--	<ul style="list-style-type: none"> • There is possibility of increase in temperature hence Provide clean, hygienic and plenty amount of drinking water to farm animals • To protect animals from heat, sprinkle cold water on animals during the afternoon, it will help to maintain the body temperature. • To reduce the stress of heat in farm animals, provide roughages by mixing with solution of 1% gaggery and 0.5% salt separately. • There is forecast for increase in temperature, hence protect animals from heat by covering roof of the shed with insulating materials such as paddy straw, dry coconut leaves and make arrangement for sprinkle cold water on the roof of shed during afternoon time. Use wet gunny bags as side curtains to protect animals from direct hot winds. • Vaccination against Haemorrhagic Septecemia (H.S) disease in farm animals under supervision of veterinary officers is advocated.
Poultry	-	<ul style="list-style-type: none"> • There is possibility of increase in temperature, hence increase the water pot and provide adequate and clean water for drinking. Also, feed should be given in the morning or evening hours. • There is forecast for increase in temperature, hence protect poultry birds from heat by covering roof of the shed with insulating materials such as paddy straw, dry coconut leaves and make arrangement for sprinkle cold water on the roof of shed during afternoon time. Use wet gunny bags as side curtains to protect poultry birds from direct hot winds. • Vaccination against Ranikhet disease in poultry birds under supervision of veterinary officers is advocated.
<p>This Agro Advisory Bulletin (AAB) is prepared and published with the consultation and recommendation of SMS committees of "Gramin Krishi Mausam Sewa (GKMS)" Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli. For more information contact nearby SAU research station or Agriculture officers of Agriculture Department, Maharashtra state.</p>		